

CD44 (Hyaluron Binding Region) M591

Mouse Monoclonal IgG1

Cat. # CM5911


Size 100 µl

Background


Cell surface adhesion protein CD44 is a ubiquitously expressed type I transmembrane protein that has important functions related to cell-cell adhesion and extracellular matrix interactions. The transmembrane protein is post-translationally modified at multiple sites by glycosylation and phosphorylation. CD44 ligands include hyaluronic acid, collagens, laminins, osteopontin, serglycin, and fibronectin. CD44 has been implicated in inflammatory cell functions as well as in tumor growth and metastasis. CD44 is overexpressed in many types of cancer; the interaction between CD44 and HER2 has been linked to an increase in ovarian carcinoma cell growth. CD44 interacts with ezrin, radixin, and moesin to link the actin cytoskeleton to the plasma membrane and the extracellular matrix. These interactions are critical for CD44 function in cell-cell adhesion and cell motility.

Background References

Tsukita, S., et al. (1994) *J Cell Biol.* 126(2):391
Goodinson, S., et al. (1999) *Mol Path.* 52(4):189
Cichy, J., et al. (2003) *J Cell Biol.* 161(5):839



Immunocytochemical labeling of CD44 in paraformaldehyde fixed human MCF7 cells. The cells were labeled with mouse monoclonal anti-CD44 (clone M591). The antibody was detected using goat anti-mouse DyLight® 594.



Western blot analysis of denatured (lanes 1 & 3) and native (lanes 2 & 4) human MDA-MB-231 whole cell lysates. The blots were probed with mouse monoclonal anti-CD44 (CM5881) at 1:1000 (lanes 1 & 2) or mouse monoclonal anti-CD44 (CM5911) at 1:1000 (lanes 3 & 4).

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

www.ecmbiosciences.com
telephone: 859-879-2075
toll-free: 1-800-859-8202
info@ecmbiosciences.com

ECMBiosciences

Rev10/16/2017

CD44 (Hyaluron Binding Region) M591

Mouse Monoclonal IgG1

Cat. # CM5911

Size 100 µl

Immunogen

Clone M591 was generated from a proprietary antigen related to the hyaluron binding region of human CD44.

Buffer and Storage

Mouse monoclonal antibody purified with protein G chromatography is supplied in 100 µl phosphate-buffered saline, 50% glycerol, 1 mg/ml BSA, and 0.05% sodium azide. Store at -20°C. Stable for 1 year.

Applications

WB	1:1000
ELISA	1:2000
ICC	1:100
IP	1:100

Species Reactivity

Hu

End user should determine optimal dilution for their particular applications and experiments.

Western blot membranes were incubated with diluted antibody in 5% non-fat milk, Tris buffer, 0.04% Tween20 for 1 hour at room temperature.

Abbreviations: E = ELISA, ICC = immunocytochemistry, IHC = immunohistochemistry, IP = immunoprecipitation, MS = mass spectrometry, WB = western blot
Hu = Human, Ms = Mouse, Rt = Rat, Ck = Chicken, F = Frog, B = Bovine

Specificity

Clone M591 detects 80-130 kDa* bands corresponding to the molecular mass of CD44 on SDS-PAGE immunoblots of denatured and native MDA-MB-231 cell lysates. The antibody also detects the denatured and native form of a recombinant human CD44 protein that includes only the hyaluron binding region. Clone M591 can be used in multiple applications including western blot, immunocytochemical labeling, ELISA, and immunoprecipitation

*All molecular weights (MW) are confirmed by comparison to MW standards and to western blot mobilities of known proteins with similar MW.

"Native" western blot utilizes non-reducing sample buffer (no mercaptoethanol or SDS), normal SDS-PAGE gel electrophoresis, and no methanol in transfer buffers.

Related Products

CM5881 CD44 (Hyaluron Binding Region) M588 Mouse Monoclonal

OM5741 Osteopontin (N-terminal region) M574 Mouse Monoclonal

IK6270 Integrin β4 Phospho-Regulation Antibody Sampler Kit

CM0071 CD44 (Extracellular region) M007 Mouse Monoclonal

CM0101 CD44 (Extracellular region) M010 Mouse Monoclonal

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

www.ecmbiosciences.com
telephone: 859-879-2075
toll-free: 1-800-859-8202
info@ecmbiosciences.com

ECMBiosciences

Rev10/16/2017