

PKC γ (C-terminus) M599

Mouse Monoclonal IgG1

Cat. # PM5991

Size 100 μ l


Background

The Protein Kinase C (PKC) family of homologous serine/threonine protein kinases is involved in a number of processes such as growth, differentiation, and cytokine secretion. At least eleven isozymes have been described. PKC consists of a single polypeptide chain containing four conserved regions (C) and five variable regions (V). The N-terminal half interacts with PKC activators Ca²⁺, phospholipid, diacylglycerol, or phorbol ester, while the C-terminal half contains the catalytic domain. The conventional PKC subfamily (α , β 1, β II, and γ) is regulated by both Ca²⁺ and diacylglycerol. The PKC pathway represents a major signal transduction system that is activated following ligand-stimulation of transmembrane receptors by hormones, neurotransmitters and growth factors. The phosphorylation of multiple sites in conventional PKCs regulates their activity. In mast cells, Fc ϵ RI stimulation leads to phosphorylation of tyrosine 658 and 662 of PKC α and PKC β I respectively. This phosphorylation requires autophosphorylation of serine 657 and 661 in these respective kinases.

Background References

Nishizuka, Y. (1988) Nature 334:661.

Kawakami et al. (2003) PNAS. USA 100:9470-9475.



Western blot analysis of PKC γ in adult mouse brain lysate. The blot was probed with mouse monoclonal anti-PKC γ clone M599 at 1:500 (lane 1) and 1:2000 (lane 2).

Applications

WB 1:1000
ELISA 1:2000

Species Reactivity

Hu, Rt, Ms, Ck

Specificity

This antibody detects an 80 kDa* protein corresponding to the molecular mass of PKC γ on SDS-PAGE immunoblots of mouse brain lysates. The antibody also detects a human recombinant PKC γ protein, but does not cross react with full length human recombinant PKC α or PKC β .

End user should determine optimal dilution for their particular applications and experiments.
Western blot membranes were incubated with diluted antibody in 5% non-fat milk, PBS, 0.04% Tween20 for 1 hour at room temperature.

*All molecular weights (MW) are confirmed by comparison to Bio-Rad Rainbow Markers and to western blot mobilities of known proteins with similar MW.

Immunogen

Clone (M599) was generated from a peptide corresponding to amino acids in the C-terminus of PKC γ . This sequence has low homology to all other PKC family members.

Buffer and Storage

Mouse monoclonal antibody purified with protein A chromatography is supplied in 100 μ l phosphate-buffered saline, 50% glycerol, 1 mg/ml BSA, and 0.05% sodium azide. Store at -20°C. Stable for 1 year.

Related Products

PM2371 PKC α (Central region) Mouse Monoclonal
PM1101 PKC (α , β 2, γ) M110 Mouse Monoclonal
PM4991 PKC (α , β , γ) M499 Mouse Monoclonal
PM5981 PKC β (N-terminal region) M598 Mouse Monoclonal
PM2421 PKC δ (N-terminal region) Mouse Monoclonal

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

www.ecmbiosciences.com
telephone: 859-879-2075
toll-free: 1-800-859-8202
info@ecmbiosciences.com

ECMBiosciences

Rev 3/29/2017