

TRPM7 (Extracellular region) M573

Mouse Monoclonal IgG1

Cat. # TM5731


Size 100 µl

Background

The transient receptor potential melastatin (TRPM) subfamily of cation-permeable TRP channels is ubiquitously expressed in mammalian tissues. This family includes TRPM1-8. In addition to acting as a calcium-permeant channel, TRPM6 and TRPM7 possess an inherent serine/threonine kinase activity. TRPM7 specifically is involved with cellular magnesium homeostasis and neurotransmitter release. Due to the magnesium inhibition, TRPM7's ion channel activity is very low. TRPM7 has been implicated in cell proliferation and migration during cancer progression, and its expression levels correlate with prognosis in breast cancer. TRPM7 kinase activation leads to massive autophosphorylation of the C-terminal region, including phosphorylation of Ser-1493, Ser-1513, and Ser-1569. Both Ser-1513 and Ser-1569 phosphorylation is required for kinase activity, and phosphorylation of Ser-1513 may inhibit Caspase-mediated cleavage of the C-terminal tail. Thus, TRPM7 is a multifunctional transmembrane protein with roles in cell signaling, proliferation, migration, and death.

Background References

Masayuki, M. et al. (2005) J of Bio Chem 280(21): 20793
Clark, K. et al. (2008) PLoS ONE 3(3): e1876
Desai, BN. et al. (2012) Dev. Cell. 22(6): 1149



Applications

WB 1:500
ELISA 1:2000

Species Reactivity

Hu, Rt, Ms

Specificity

The antibody was affinity purified using TRPM7 (Extracellular region) peptide. This antibody detects a 220 kDa* protein on SDS-PAGE immunoblots of human MCF7 cells.

End user should determine optimal dilution for their particular applications and experiments.
Western blot membranes were incubated with diluted antibody in 5% non-fat milk, PBS, 0.04% Tween20 for 1hour at room temperature.

*All molecular weights (MW) are confirmed by comparison to Bio-Rad Rainbow Markers and to western blot mobilities of known proteins with similar MW.

Immunogen

TRPM7 synthetic peptide (coupled to carrier) corresponding to amino acids in the extracellular region of human TRPM7. This site is well conserved in rat and mouse TRPM7, but has low homology to other TRPM family members.

Buffer and Storage

Mouse monoclonal, affinity-purified antibody is supplied in 100µl phosphate-buffered saline, 50% glycerol, 1 mg/ml BSA, and 0.05% sodium azide. Store at -20°C. Stable for 1 year.

Related Products

TP5651 TRPM7 (Extracellular region) Rabbit Polyclonal
TP5691 TRPM7 (a.a. 1484-1497) Rabbit Polyclonal
TP5661 TRPM7 (Ser-1493), phospho-specific Rabbit Polyclonal
TP5671 TRPM7 (Ser-1513), phospho-specific Rabbit Polyclonal
TP5681 TRPM7 (Ser-1569), phospho-specific Rabbit Polyclonal

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

www.ecmbiosciences.com
telephone: 859-879-2075
toll-free: 1-800-859-8202
info@ecmbiosciences.com

ECMBiosciences

Rev 6/29/2016