

# TRPM7 (Ser-1493), phospho-specific Rabbit Polyclonal

Cat. # TP5661


Size 100 µl

## Background

The transient receptor potential melastatin (TRPM) subfamily of cation-permeable TRP channels is ubiquitously expressed in mammalian tissues. This family includes TRPM1-8. In addition to acting as a calcium-permeant channel, TRPM6 and TRPM7 possess an inherent serine/threonine kinase activity. TRPM7 specifically is involved with cellular magnesium homeostasis and neurotransmitter release. Due to the magnesium inhibition, TRPM7's ion channel activity is very low. TRPM7 has been implicated in cell proliferation and migration during cancer progression, and its expression levels correlate with prognosis in breast cancer. TRPM7 kinase activation leads to massive autophosphorylation of the C-terminal region, including phosphorylation of Ser-1493, Ser-1513, and Ser-1569. Both Ser-1513 and Ser-1569 phosphorylation is required for kinase activity, and phosphorylation of Ser-1513 may inhibit Caspase-mediated cleavage of the C-terminal tail. Thus, TRPM7 is a multifunctional transmembrane protein with roles in cell signaling, proliferation, migration, and death.

## Background References

Masayuki, M. et al. (2005) J of Bio Chem 280(21): 20793  
Clark, K. et al. (2008) PLoS ONE 3(3): e1876  
Desai, BN. et al. (2012) Dev. Cell. 22(6): 1149



## Applications

WB 1:500  
ELISA 1:1000

## Species Reactivity

Hu

## Specificity

The antibody was cross-adsorbed to unphosphorylated TRPM7 (Ser-1493) then affinity purified using phospho-TRPM7 (Ser-1493) peptide. This antibody detects a 220 kDa\* protein on SDS-PAGE immunoblots of human MDA-MB-231 cells stimulated with calyculin A, and detects a 70 kDa autophosphorylated recombinant kinase domain of human TRPM7. These reactivities are not observed after lambda phosphatase treatment to dephosphorylate TRPM7.

\*All molecular weights (MW) are confirmed by comparison to Bio-Rad Rainbow Markers and to western blot mobilities of known proteins with similar MW.

## Immunogen

Phospho-TRPM7 (Ser-1493) synthetic peptide (coupled to carrier) corresponding to amino acids surrounding Ser-1493 in human TRPM7. This site is conserved in rat and mouse TRPM7, but the homology surrounding the site is low. This site has no homology to other TRPM family members.

## Buffer and Storage

Rabbit polyclonal, affinity-purified antibody is supplied in 100µl phosphate-buffered saline, 50% glycerol, 1 mg/ml BSA, and 0.05% sodium azide. Store at -20°C. Stable for 1 year.

## Related Products

TP5651 TRPM7 (Extracellular region) Rabbit Polyclonal  
TP5691 TRPM7 (a.a. 1484-1497) Rabbit Polyclonal  
TP5671 TRPM7 (Ser-1513), phospho-specific Rabbit Polyclonal  
TP5681 TRPM7 (Ser-1569), phospho-specific Rabbit Polyclonal  
TP5701 TRPM8 (Extracellular region) Rabbit Polyclonal

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

www.ecmbiosciences.com  
telephone: 859-879-2075  
toll-free: 1-800-859-8202  
info@ecmbiosciences.com

**ECM**Biosciences

Rev 7/11/2017